

ARMY RESEARCH LABORATORY



# Scavenging Models for Smokes

Steven M. Kovel

ARL-MR-503

February 2001

Approved for public release; distribution unlimited.

20010326 037

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

# Army Research Laboratory

Adelphi, MD 20783-1197

---

ARL-MR-503

February 2001

---

## Scavenging Models for Smokes

Steven M. Kovel

Computational and Information Sciences Directorate

---

Approved for public release; distribution unlimited.

---

---

## Abstract

---

The smoke and obscuration computer model known as the Combined Obscuration Model for Battlefield Induced Contaminants (COMBIC) has the potential for treating the effect of precipitation scavenging as one of the model's input parameters. This report examines the impact of rainfall on the evolution of a white phosphorous (WP) smoke cloud based upon the predictions of the model. The result of the analysis indicates that the effect of precipitation scavenging is less than the error expected in COMBIC predictions. Therefore, the recommendation is not to incorporate more elaborate precipitation models in COMBIC.

---

## Contents

---

1	Introduction	1
2	Scavenging	2
3	Number Distributions	4
3.1	White Phosphorous Distribution . . . . .	4
3.2	Raindrop Distribution Function . . . . .	5
4	Cloud Growth Modeling	7
5	Conclusions	10
	References	11
	Distribution	13
	Report Documentation Page	17

## Figures

1	Scavenging coefficient versus aerosol particle size . . . . .	3
2	White phosphorous distribution . . . . .	5
3	Raindrop distribution . . . . .	5

## Tables

1	Comparison of two scavenging values . . . . .	8
1	Comparison of two scavenging values (cont'd) . . . . .	9

---

## 1. Introduction

---

The study of precipitation scavenging has an extensive history and development that began with the desire to model clouds and rain. The term "precipitation scavenging" means the removal of material present in the atmosphere by falling precipitation. When clouds form, the atmospheric water vapor is collected around a nucleation particle that floats within the cloud. As the particle moves, additional water is collected until the particle reaches sufficient size and mass to begin falling to the ground. This description and representation of the phenomenology can be traced back to the analysis of the operation of the Wilson cloud chamber (which was in use in 1903), if not earlier. Currently, the description has been extended to include many additional scientific representations including such complex phenomena as the turbulent motion of aerosols that occur within a storm cloud and the chemical interaction of the particles.

Scavenging also takes place when the precipitate from the cloud passes through aerosols nearer to the ground. Work on this phenomenology began to attract attention when a national concern about nuclear fallout was expressed. This topic received an additional analytical impetus in the 1960s and 1970s when environmental pollution began to achieve national importance. Many sophisticated models have evolved from this extensive work as well as some simple parametric representations of the phenomenology. This report addresses the question as to whether an appropriate choice for the model can be identified that can be integrated with the Combined Obscuration Model for Battlefield Induced Contaminants (COMBIC) [1]. This model predicts the growth and extent of battlefield obscurants as well as the transmission through the obscurant. Knowing the extent to which the obscurant could be reduced by precipitation scavenging could be tactically significant.

---

## 2. Scavenging

---

The theoretical development of many scavenging models is formulated from material balance considerations [2]. Alternate approaches to this development can be based either on energy considerations or momentum considerations. A general form of the differential equation representing the material balance for the change of the aerosol concentration is given by the following equation:

$$\frac{\partial n}{\partial t} = -\nabla \cdot (n \bar{v}) + w, \quad (1)$$

where  $\bar{v}$  is the velocity of the aerosol and  $w$  is a source term. Note that all the terms in this equation can be functions of both the position,  $r$ , and time,  $t$ . In this analysis we will take the source term to be zero.

Following the analysis presented by Pruppacher and Klett [3, p. 380, eq 12-80], the scavenging coefficient,  $\Lambda$ , is defined by the equation

$$-\frac{1}{n} \frac{\partial n(r, t)}{\partial t} \cong \Lambda(r, t), \quad (2)$$

where  $n(r, t)$  is the aerosol concentration. A very simple model occurs if the scavenging coefficient does not vary with time. Then the last relationship can be integrated and written as

$$n(r, t) = n(r, 0)e^{-\Lambda(r)t}. \quad (3)$$


Within COMBIC, a modified form of this equation is used to model the process for removing mass by deposition on the ground or evaporation. A constant additional parameter,  $f_d$ , is added to the equation. The purpose of this parameter is to represent a residue remaining after removing all that can possibly be scavenged. When this parameter is included, the equation takes the form

$$n(r, t) = n(r, 0)[f_d + (1 - f_d)e^{-\Lambda(r)t}]. \quad (4)$$

However, within the context of this analysis, this parameter should be set to zero.

The value for the scavenging coefficient depends on the scavenging mechanism prevalent for the aerosol particle size being considered. For a small aerosol particle with a radius less than  $0.1 \mu\text{m}$ , the primary capture mechanism is Brownian diffusion. For particles with a radius greater than  $1.0 \mu\text{m}$ , the primary capture mechanism is inertial impaction. Particles with radius values falling between these two values (commonly called the Greenfield gap) have a number of other mechanisms that generate the scavenging. Figure 1 shows a curve of the scavenging coefficient values as a function of aerosol particle size for two different distributions of raindrop size [3, p. 395, fig. 12-10].

Figure 1. Scavenging coefficient versus aerosol particle size [3].



In this figure, the term phoretic effect refers to two phenomena called thermophoresis and diffusiophoresis. The former refers to the motion of particles arising from the nonuniform heating of particles due to temperature gradients in the suspending gas. Diffusiophoresis refers to particle motion arising from concentration gradients in a gaseous mixture. The parameter  $a_{\text{max}}$  is a value that determines the maximum raindrop size for the distribution function. This distribution function will be discussed below. Other parameters include the temperature difference between the surface of a droplet and the bulk aerosol of  $3^\circ\text{C}$ , a precipitation rate of  $10 \text{ mm hr}^{-1}$ , and a drop terminal velocity given by the relationship  $(8000 a) \text{ cm s}^{-1}$ , with  $a$  in cm. Note that in these curves, a particle radius of greater than  $\sim 0.5 \mu\text{m}$  or less than  $\sim 0.002 \mu\text{m}$  would be required to obtain a scavenging coefficient greater than  $0.1 \text{ hr}^{-1}$ . The value of  $0.1 \text{ hr}^{-1}$  is relatively small, since it represents only a 10-percent decrease in the particle density after 1 hr of precipitation.



---

### 3. Number Distributions

---

In the following analysis, I focused on white phosphorous (WP) as the obscurant and wanted to select a single value for the WP particle radius and a single value for the raindrop size. This does not represent a real situation. Each of the particles has an associated size distribution and I need to examine how well a single value represents a realistic situation.

#### 3.1 White Phosphorous Distribution

According to the information in [4], experiments by Jennings and Gillespie in 1978 [5] have shown that the particle size spectrum is closely approximated by a lognormal distribution [4, p. 31]. This distribution is given to be [4, p. 18, eq 19]

$$n(r) = \left[ \frac{N}{r\sqrt{2\pi \ln(\sigma_g)}} \right] \exp \left[ \frac{-(\ln r - \ln r_g)^2}{2(\ln(\sigma_g))^2} \right], \quad (5)$$

where

$\sigma_g$  = geometric mean standard deviation,  
 $r_g$  = geometric mean radius in  $\mu\text{m}$ , and  
 $N$  = aerosol particle number density (particles per  $\text{cm}^3$ ).

Another equation given in the report [4, p. 19, eq 21] relates these parameters to the mass concentration  $C$  by the relationship

$$C = \frac{4}{3}\pi\rho N r_g^3 \exp \left[ \frac{9}{2}(\ln \sigma_g)^2 \right]. \quad (6)$$

Values for these parameters are given in [4, p. 32, table 16]. Since I am interested in precipitation scavenging, the values I take here are for WP at a 90-percent relative humidity. For that case

$r_g = 0.365 \mu\text{m}$ ,  
 $\sigma_g = 1.450$ ,  
 $\rho = 1.178 \text{ g/cm}^3$ , and  
 $C = 10^6 \mu\text{g/m}^3$ .

Note that the value used for  $C$  is a representative value that was appropriate for the analysis being done in the report [4]. Since I am concerned only with the relative distribution of particle sizes, that value will be used here. However, before making the substitution, I obtain the relationship between the mass density and the number density. Substituting these values into the last relationship yields

$$C = 4.466 * 10^{-13} * N. \quad (7)$$

For the representative mass-loading value of  $10^6 \mu\text{g}/\text{m}^3$ , then, the particle density  $N$  is determined to be  $2.24 \times 10^6$  particles/ $\text{cm}^3$ .

Figure 2 shows a plot of the function for these values. Note that the particle density is not a sharp function of the particle radius. For the number density to drop by a factor of two from its peak value at 0.365 micrometers, the radius would have to be either as small as about 0.2  $\mu\text{m}$  or as large as about 0.5  $\mu\text{m}$ . While this particle size range lies within the inertial impaction portion of the scavenging coefficient curve, it does extend into the Greenfield gap portion of that curve.

### 3.2 Raindrop Distribution Function

The raindrop size distribution function used in the preceding section is given by the relationship (see the caption for figure 12- 10 in [3]),

$$n(a) = 10^{-4} \frac{R}{6} \pi a_{\max}^7 a^2 \exp\left(-2 \frac{a}{a_{\max}}\right), \quad (8)$$

where  $a$  is the raindrop size in cm and  $R$  is the precipitation rate in  $\text{cm s}^{-1}$ . This relationship is also known as the Khrgian-Mazin drop-size distribution. See figure 3 for a plot of these distributions. The value used for the precipitation rate  $R$  in the curves is equal to  $1/3600 \text{ cm s}^{-1} = 10 \text{ mm/hr}$ . In the PFNDAT report [4, p. 24, table 9], a rain rate of 10 mm/hr is termed a thunderstorm, 5 mm/hr is a widespread rain, and 1 mm/hr is a drizzle.

Depending on other environmental conditions such as vertical wind draft, different values can be obtained for the maximum value size. No rationale

Figure 2. White phosphorous distribution.





Figure 3. Raindrop distribution.



for selecting one size over another exists. This implies that we will want to consider the full dynamic range of the scavenging coefficient that is restricted only by the WP particle size. Based on the scavenging curve, the range of values for the scavenging coefficient  $\Lambda$  will range from about  $2 \times 10^{-4}$  per hr to  $5 \times 10^{-2}$  per hr. The impact of this range of values on the growth of a WP smoke cloud will be examined in the next section.

---

## 4. Cloud Growth Modeling

---

The model to be used to examine the growth of a WP smoke cloud is COMBIC [1], which has two phases for computing the growth of battle-field obscurants such as dust, smoke, and clouds, and for determining atmospheric extinction along user-determined line-of-sight. The model has evolved through several versions beginning in 1982 to 1992. Source code is written in FORTRAN 77.

COMBIC can be separated into two parts. The first part computes the growth of a puff or a plume under the influence of diffusion, gravity, and external winds. A basic assumption for the computation is that the cloud is an ellipsoidal Gaussian distribution. The physical equations that describe the growth will maintain the Gaussian form of the result, although the parameters describing the shape will deform. Results of this computation are placed into an auxiliary database that is indexed based on the time since the cloud was started. If several different types of clouds are generated, each one is accorded its own descriptive growth. This database is available for further analysis and is the basis for the following discussion.

Two important points with regard to COMBIC need to be highlighted: first, an error exists in some versions of COMBIC in the way parameter initialization is performed when a nonzero scavenging coefficient is used. This problem has been corrected in the code used here. However, if a reader using the software has scavenging results that are peculiar, the code used to duplicate these results may not reflect the required correction. Second, a zero wind speed should not be used since it can cause a discontinuity in the computation for cloud growth. In part of the code, a minimum value of 0.25 m/s is required and is automatically set for the computation. An undocumented recommendation is that the lowest wind speed that should be used is 1 m/s.

Two runs were made of COMBIC. The first run assumed that no scavenging was taking place. The second run assumed that a scavenging coefficient of  $2 \times 10^{-5} \text{ s}^{-1}$  was appropriate for representing the WP distribution. All other environmental parameters were the same (wind speed = 1 m/s, relative humidity 90 percent). The results of these computations are given in table 1.

Note that the maximum mass difference between the scavenging and nonscavenging cases does not exceed 1.5 percent of the total mass produced. Similarly, the transmission measured between two points perpendicular to the windborne flow of the smoke does not show any significant differences between the two cases. The wind speed has a greater effect on the amount of time the obscurant remains within a given location, reducing

the transmission. Within the accuracy of this model, scavenging effects can be ignored for determining the airborne mass and transmission of WP.

A quick check was run with the use of a scavenging coefficient that was 10 times larger and then again with the coefficient 100 times larger. Only in the latter case did significant effects begin to occur. Since those values are completely outside the range of the data considered here, the results were ignored.

Table 1. Comparison of two scavenging values.

Time (s)	Mass produced	Airborne $\Lambda = 0$	Airborne $\Lambda = .00002$	% Difference/ 100
1	189.6	189.6	189.6	0.000
10	1865.5	1865.5	1865.1	0.000
20	3663.4	3663.4	3661.9	0.000
30	5395.4	5395.4	5392.2	0.001
40	7063.5	7063.5	6995.9	0.010
50	8669.1	8669.1	8590.5	0.009
60	10214.1	10214.1	10122.8	0.009
70	11700.2	11700.2	11594.3	0.009
80	13128.9	13128.9	13011.8	0.009
90	14502.0	14502.0	14376.8	0.009
100	15821.0	15821.0	15688.2	0.008
110	17087.5	17087.5	16947.9	0.008
120	18303.1	18303.1	18157.4	0.008
130	19469.4	19469.4	19317.5	0.008
140	20587.7	20587.7	20430.0	0.008
150	21659.7	21659.7	21496.1	0.008
160	22686.8	22686.8	22517.2	0.007
170	23670.5	23670.5	23494.8	0.007
180	24612.2	24612.2	24430.3	0.007
190	25513.2	25513.2	25325.1	0.007
200	26375.1	26375.1	26180.5	0.007
210	27199.0	27199.0	26998.0	0.007
220	27986.4	27986.4	27778.7	0.007
230	28738.6	28738.6	28524.2	0.007
240	29456.9	29456.9	29235.6	0.008
250	30142.5	30142.5	29914.3	0.008
260	30796.7	30796.7	30561.4	0.008
270	31420.7	31420.7	31178.3	0.008
280	32015.7	32015.7	31766.1	0.008
290	32582.8	32582.8	32326.1	0.008
300	33123.3	33123.3	32859.3	0.008
310	33638.2	33638.2	33366.9	0.008
320	34128.8	34128.8	33850.1	0.008
330	34595.9	34595.9	34309.9	0.008
340	35040.8	35040.8	34747.3	0.008
350	35464.4	35464.4	35163.5	0.008
360	35867.8	35867.8	35559.4	0.009
370	36251.9	36251.9	35936.0	0.009
380	36617.7	36617.7	36294.4	0.009
390	36966.3	36966.3	36635.4	0.009
400	37298.4	37298.4	36960.0	0.009
410	37615.0	37615.0	37269.1	0.009
420	37917.0	37917.0	37563.5	0.009
430	38205.3	38205.3	37844.2	0.009
440	38480.6	38480.6	38111.9	0.010

Table 1. Comparison of  
two scavenging values  
(cont'd).

Time (s)	Mass produced	Airborne $\Lambda = 0$	Airborne $\Lambda = .00002$	% Difference/ 100
450	38743.7	38743.7	38367.4	0.010
460	38995.6	38995.6	38611.5	0.010
470	39236.8	39236.8	38844.8	0.010
480	39468.3	39468.3	39067.7	0.010
490	39690.2	39690.2	39280.7	0.010
500	39902.7	39902.7	39484.3	0.010
510	40106.2	40106.2	39678.7	0.011
520	40301.1	40301.1	39864.6	0.011
530	40487.7	40487.7	40042.1	0.011
540	40666.4	40666.4	40211.6	0.011
550	40837.5	40837.5	40373.6	0.011
560	41001.3	41001.3	40528.3	0.012
570	41158.2	41158.2	40676.0	0.012
580	41308.5	41308.5	40817.1	0.012
590	41452.4	41452.4	40951.7	0.012
600	41590.2	41590.2	41080.3	0.012
610	41722.1	41722.1	41203.0	0.012
620	41848.4	41848.4	41320.1	0.013
630	41969.4	41969.4	41431.8	0.013
640	42085.3	42085.3	41538.4	0.013
650	42196.2	42196.2	41640.1	0.013
660	42302.4	42302.4	41737.1	0.013
670	42404.2	42404.2	41829.6	0.014
680	42501.6	42501.6	41917.8	0.014
690	42594.9	42594.9	42001.8	0.014
700	42684.2	42684.2	42081.9	0.014
710	42769.7	42769.7	42158.2	0.014
720	42851.7	42851.7	42230.8	0.014
730	42930.1	42930.1	42300.1	0.015
740	43005.2	43005.2	42365.9	0.015
750	43077.1	43077.1	42435.8	0.015
760	43146.0	43146.0	42495.2	0.015
770	43212.0	43212.0	42551.6	0.015
780	43275.1	43275.1	42605.3	0.015

---

## 5. Conclusions

---

This simplified model predicts that the effect of scavenging on a WP smoke is minimal. While some slight changes occur both in the mass produced by the burning source and the transmission through the resulting cloud, the differences between these numbers are not great enough to warrant confidence when using results that cover long periods of time. In addition, since the assumptions about both raindrop and smoke particle size had the effect of maximizing the scavenging coefficient, the error would be on the side of a worst-case analysis, that is, the true effect of scavenging is even less than these results would indicate.

From a tactical viewpoint, when small amounts of smoke are deployed, it is better to wait for the wind to remove the smoke than depend on the scavenging by precipitation. For large amounts of smoke that are continuously deployed, scavenging is not an important factor.

I made a search to determine if any experimental data supporting the scavenging coefficient used for WP exist. None were found. The major reason for this omission is that the equipment used when WP and other militarily significant smokes were measured would not operate in a rainy environment. Measurements on WP are available for fair weather and snowy conditions.

Consider the original question of whether an appropriate choice for the scavenging model can be identified that can be combined with COMBIC. A simple model is already present within COMBIC requiring the input of only a single parameter. Until experimental data can be obtained demonstrating an error in the estimates obtained with the use of this simple approach, enhancing the representation is not necessary. Of greater importance are the effects of terrain and nonuniform wind fields on the scavenging of smokes and other obscurants.

---

## References

---

1. Wetmore, A., and S. D. Ayres (August 2000), "COMBIC, Combined Obscuration Model for Battlefield Induced Contaminants," 2 vols, U.S. Army Research Laboratory, ARL-TR-1831-1, -2.
2. Hales, J. M. (1986), "The Mathematical Characterization of Precipitation Scavenging and Precipitation Chemistry," *The Handbook of Environmental Chemistry*, vol 4, part A.
3. Pruppacher, H. R., and J. D. Klett (1984), *Microphysics of Clouds and Precipitation*, D. Reidel Publishing Company.
4. Tofsted, D. H., B. T. Davis, A. E. Wetmore, J. Fitzgerrel, R. C. Shirkey, and R. A. Sutherland (June 1997), *EOSAEL92, Aerosol Phase Function Data Base PFNDAT*, U.S. Army Research Laboratory, ARL-TR-273-9.
5. Jennings, S. G., and J. B. Gillespie (1978), *Mie Theory Sensitivity Studies—The Effects of Aerosol Complex Refractive Index and Size Distribution Variations on Extinction and Absorption Coefficients*, U. S. Army Atmospheric Sciences Laboratory, ASL-TR-0003, White Sands Missile Range, NM.



## Distribution

Admnstr  
Defns Techl Info Ctr  
ATTN DTIC-OCF  
8725 John J Kingman Rd Ste 0944  
FT Belvoir VA 22060-6218

DARPA  
ATTN S Welby  
3701 N Fairfax Dr  
Arlington VA 22203-1714

Mil Asst for Env Sci Ofc of the Undersec of  
Defns for Rsrch & Engrg R&AT E LS  
Pentagon Rm 3D129  
Washington DC 20301-3080

Ofc of the Secy of Defns  
ATTN ODDRE (R&AT)  
The Pentagon  
Washington DC 20301-3080

Ofc of the Secy of Defns  
ATTN OUSD(A&T)/ODDR&E(R) R J Trew  
3080 Defense Pentagon  
Washington DC 20301-7100

AMCOM MRDEC  
ATTN AMSMI-RD W C McCorkle  
Redstone Arsenal AL 35898-5240

ARL Chemical Biology Nuc Effects Div  
ATTN AMSRL-SL-CO  
Aberdeen Proving Ground MD 21005-5423

Army Corps of Engrs Engr Topographics Lab  
ATTN CETEC-TR-G P F Krause  
7701 Telegraph Rd  
Alexandria VA 22315-3864

Army Field Artillery Schl  
ATTN ATSF-TSM-TA  
FT Sill OK 73503-5000

Army Infantry  
ATTN ATSH-CD-CS-OR E Dutoit  
FT Benning GA 30905-5090

US Army TRADOC  
Battle Lab Integration & Techl Dirctr  
ATTN ATCD-B J A Klevecz  
FT Monroe VA 23651-5850

US Military Acdmly  
Mathematical Sci Ctr of Excellence  
ATTN MADN-MATH MAJ M Huber  
Thayer Hall  
West Point NY 10996-1786

Natl Ground Intllgnc Ctr Army Foreign Sci  
Tech Ctr  
ATTN CM  
220 7th Stret NE  
Charlottesville VA 22901-5396

Natl Security Agcy  
ATTN W21 Longbothum  
9800 Savage Rd  
FT George G Meade MD 20755-6000

SMC/CZA  
2435 Vela Way Ste 1613  
El Segundo CA 90245-5500

US Army CRREL  
ATTN CRREL-GP R Detsch  
ATTN Roberts  
ATTN SWOE G Koenig  
72 Lyme Rd  
Hanover NH 03755-1290

US Army Info Sys Engrg Cmnd  
ATTN AMSEL-IE-TD F Jenia  
FT Huachuca AZ 85613-5300

US Army Natl Ground Intllgnc Ctr  
ATTN IANG-TSC J Breeden  
220 Seventh Stret NE  
Charlottesville VA 22902

US Army OEC  
ATTN CSTE-AEC-FSE  
4501 Ford Ave Park Center IV  
Alexandria VA 22302-1458

US Army Simulation Train & Instrmntn  
Cmnd  
ATTN J Stahl  
12350 Research Parkway  
Orlando FL 32826-3726

## Distribution (cont'd)

US Army STRICOM  
ATTN AMCPM-WARSIM Lavin  
ATTN AMCPM-WARSIM S Veautour  
ATTN AMCPM-WARSIM Williams  
Orlando FL 32826

US Army Tank-Automtv Cmnd RDEC  
ATTN AMSTA-TR J Chapin  
Warren MI 48397-5000

US Army Topo Engrg Ctr  
ATTN CETEC-ZC  
FT Belvoir VA 22060-5546

US Army TRADOC Anal Cmnd-WSMR  
ATTN ATRC-WSS-R  
White Sands Missile Range NM 88002

US Army White Sands Missile Range  
ATTN STEWS-IM-ITZ Techl Lib Br  
White Sands Missile Range NM 88002-5501

US Army TRADOC  
ATTN ATRC-WEC D Dixon  
White Sands Missile Range NM 88002-5501

USATRADOC  
ATTN ATCD-FA  
FT Monroe VA 23651-5170

Nav Air War Cen Wpn Div  
ATTN CMD 420000D C0245 A Shlanta  
1 Admin Cir  
China Lake CA 93555-6001

Nav Air Warfare Ctr Train Sys Div  
ATTN Code 494 A Hutchinson  
12350 Research Parkway  
Orlando FL 32826

Nav Surfc Warfare Ctr  
ATTN Code B07 J Pennella  
17320 Dahlgren Rd Bldg 1470 Rm 1101  
Dahlgren VA 22448-5100

Naval Surfc Weapons Ctr  
ATTN Code G63  
Dahlgren VA 22448-5000

AFCCC/DOC  
ATTN Glauber  
151 Patton Ave Rm 120  
Asheville NC 28801-5002

AFSPC/DRFN  
ATTN CAPT R Koon  
150 Vandenberg Stret Ste 1105  
Peterson AFB CO 80914-45900

Phillips Lab  
ATTN PL/LYP  
Hanscom AFB MA 01731-5000

Phillips Lab Atmos Sci  
Div Geophysics Dirtrt  
ATTN PL-LYP Chisholm  
Kirtland AFB NM 87118-6008

USAF Rome Lab Tech  
ATTN Corridor W Ste 262 RL SUL  
26 Electr Pkwy Bldg 106  
Griffiss AFB NY 13441-4514

NASA Marshal Spc Flt Ctr  
Atmos Sci Div  
ATTN Code ED 41 1  
Huntsville AL 35812

NIST  
ATTN MS 847.5 M Weiss  
325 Broadway  
Boulder CO 80303

Hicks & Assoc Inc  
ATTN G Singley III  
1710 Goodrich Dr Ste 1300  
McLean VA 22102

Natl Ctr for Atmos Rsrch  
ATTN NCAR Library Serials  
PO Box 3000  
Boulder CO 80307-3000

NCSU  
ATTN J Davis  
PO Box 8208  
Raleigh NC 27650-8208

## Distribution (cont'd)

OL-A/AFCCC/MST  
ATTN G McWilliams  
151 Patton St Rm 120  
Asheville NC 28801-5002

Pacific Mis Test Ctr Geophysics Div  
ATTN Code 3250  
Point Mugu CA 93042-5000

U.S. Army Rsrch Lab  
ATTN AMSRL-IS-EW  
White Sands Missile Range NM 88002-5501

US Army Rsrch Lab  
ATTN BED IS&TD R Shirkey  
White Sands Missile Range NM 88002

US Army Rsrch Lab  
ATTN AMSRL-CI-EW D Hoock  
Battlefield Envir Div  
White Sands Missile Range NM 88002-5001

US Army Rsrch Lab  
ATTN AMSRL-SL J Wade  
White Sands Missile Range NM 88002

US Army Rsrch Ofc  
ATTN AMSRL-RO-EN W D Bach  
PO Box 12211  
Research Triangle Park NC 27709

US Army Rsrch Lab  
ATTN AMSRL-CI-AI-R Mail & Records  
Mgmt  
ATTN AMSRL-CI-AP Techl Pub (2 copies)  
ATTN AMSRL-CI-LL Techl Lib (2 copies)  
ATTN AMSRL-DD J M Miller  
ATTN AMSRL-CI J D Gantt  
ATTN AMSRL-CI-EP S Kovel (10 copies)  
ATTN AMSRL-SE J Pellegrino  
ATTN AMSRL-SE-EE Z G Sztankay  
Adelphi MD 20783-1197

<b>REPORT DOCUMENTATION PAGE</b>			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE February 2001		3. REPORT TYPE AND DATES COVERED Final, FY 2000
4. TITLE AND SUBTITLE Scavenging Models for Smokes			5. FUNDING NUMBERS DA PR: AH71 PE: 622784H71	
6. AUTHOR(S) Steven M. Kovel				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory Attn: AMSRL-CI-EP email: skovel@arl.army.mil Adelphi, MD 20783-1197			8. PERFORMING ORGANIZATION REPORT NUMBER ARL-MR-503	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory 2800 Powder Mill Road Adelphi, MD 20783-1197			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES ARL PR: 1FEH16 AMS code: 622784H7111				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The smoke and obscuration computer model known as the Combined Obscuration Model for Battlefield Induced Contaminants (COMBIC) has the potential for treating the effect of precipitation scavenging as one of the model's input parameters. This report examines the impact of rainfall on the evolution of a white phosphorous (WP) smoke cloud based upon the predictions of the model. The result of the analysis indicates that the effect of precipitation scavenging is less than the error expected in COMBIC predictions. Therefore, the recommendation is not to incorporate more elaborate precipitation models in COMBIC.				
14. SUBJECT TERMS Modeling, interfaces, smokes			15. NUMBER OF PAGES 20	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	